

**Title:** *Early transmissibility assessment of a novel coronavirus in Wuhan, China*

**Authors:** Maimuna S. Majumder, PhD, MPH<sup>1,2</sup> and Kenneth D. Mandl, MD, MPH<sup>1,2</sup>

**Affiliations:** <sup>1</sup>Computational Health Informatics Program, Boston Children's Hospital, Boston, MA, USA; <sup>2</sup>Department of Pediatrics, Harvard Medical School, Boston, MA, USA

**Author Information:**

Name: Maimuna S. Majumder

Phone: 1-978-460-3677

Email Address: [maimuna.majumder@childrens.harvard.edu](mailto:maimuna.majumder@childrens.harvard.edu)

Mailing Address: 300 Longwood Avenue, Boston, MA 02115

Name: Kenneth D. Mandl

Phone: 617-355-4145

Email Address: [Kenneth.Mandl@Harvard.edu](mailto:Kenneth.Mandl@Harvard.edu)

Mailing Address: 300 Longwood Avenue, Boston, MA 02115

**Manuscript Word Count:** 713

**Funding Statement:** This work was supported in part by grant T32HD040128 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## COMMENT:

Since December 1, 2019, nearly 3000 cases of respiratory illness caused by a novel coronavirus (2019-nCoV) have been diagnosed in relation to an outbreak originating in Wuhan, China [1, 2]. Though several cases have been linked to a local fish and live animal market, human-to-human transmission appears likely and cases have already been exported to neighboring countries [3]. Due to the novelty of this pathogen, rapid assessment of its associated epidemiological properties is of the essence – especially within the context of the ease with which it may be transmitted. In this Comment, we present a cumulative epidemic curve of cases that have been reported to date and assess the potential transmissibility associated with 2019-nCoV given currently available public information.

Cumulative case count data were collected from the World Health Organization Disease Outbreak News reports; bulletins issued by the Chinese National Health Commission and the Wuhan Municipal Health Commission; and a recent study published in *The Lancet* [1–4]. Upon plotting these data over time, a reverse-L-shaped curve appeared due to reporting of large case counts from January 18–26, 2020. Though not necessarily representative of true case growth, such a shape is not uncommon for novel pathogens and can be driven by early delays in case identification and reporting (Figure 1A) [5].

This cumulative epidemic curve was then used to assess potential transmissibility via the Incidence Decay and Exponential Adjustment (IDEA) model [6]. The IDEA model is a simple, phenomenological method that allows for estimation of the basic reproduction number ( $R_0$ ): the average number of individuals a new case is expected to infect in a fully susceptible population. When a cumulative epidemic curve of a disease is available and

when the length of time between two consecutive infections in a chain of transmission (i.e. the serial interval of the pathogen) can be approximated, this model can be employed in otherwise data-scarce settings. Previous studies have demonstrated performance of the IDEA model to be on-par with more traditional compartmental (e.g. S-I-R) models for multiple emerging diseases, including Zika, Ebola, and Middle East Respiratory Syndrome (MERS) [5-8].

The model itself can be defined by the following single equation:

$$I = \left[ \frac{R_0}{(1 + d)^t} \right]^t$$

Here,  $t$  is the number of serial intervals that have passed at time of model parameterization and  $I$  is incidence at serial interval  $t$ . Meanwhile,  $d$  is a discount factor that takes into account reductions in transmissibility over time due to the natural depletion of susceptible individuals in the affected population and any public interventions that may impact disease spread over time.

Because the serial interval associated with 2019-nCoV has not yet been established, we referenced mean serial interval lengths ( $I$ ) from the related SARS-Coronavirus and MERS-Coronavirus [range: 6-10 days] to parameterize our model [9, 10]. For this range of serial interval lengths, modeled  $R_0$  estimates varied from 2.0 to 3.1 when using data from December 8, 2019 through January 26, 2020. Estimates for  $d$  were 0 for all serial interval lengths. Fits are shown in Figure 1A for a mean  $R_0$  of 2.5, in which the model aims to standardize the reverse-L-shaped curve against a more realistic exponential growth curve.

To ascertain the stability of our estimates thus far, we also used truncated input data (i.e. December 8, 2019 through January 18, 19, 20, and so on, through January 26) to parameterize our model. Results for the estimated reproduction number range were similar to those obtained when using all available data, signaling relative short-term stability in our model's ability to capture the growth rate of the outbreak over the last 9 days, despite elevated case count reporting in this time frame (Figure 1B).


Because  $R_0$  values greater than 1 indicate the possibility of sustained transmission, our analysis suggests that 2019-nCoV may exhibit epidemic potential under the assumption that currently available public information is accurate and relatively complete. Notably, our current estimates of  $R_0$  – while preliminary and subject to change – are consistent with previous estimates associated with SARS-Coronavirus ( $R_0 = 2-4$ ) [9]. With this in mind, timely data sharing and further epidemiological research should be pursued actively as the ongoing outbreak progresses. Pending availability of new data, repeated re-evaluation of the model presented here – as well as the transmissibility associated with 2019-nCoV more broadly – is recommended.

## Figure Legend:


**Figure 1.** *Cumulative epidemic curve, model fits, and early transmissibility assessment for novel coronavirus cases associated with Wuhan, China from December 1, 2019 through January 26, 2020.* In Panel A (top), reported cumulative cases are shown with pink circular markers connected by a pink dotted line and model fits for a mean  $R_0$  of 2.5 are shown with a blue dotted line using data from December 1, 2019 through January 26, 2020. In Panel B (bottom), reproduction number ranges as estimated by the Incidence Decay and Exponential Adjustment (IDEA) model are shown using data through January 18, 19, 20, and so on, through January 26 to demonstrate short-term model stability as well as transmissibility. Please note that on January 24, Huang et al. published a study in which the start date of the outbreak was revised backward 7 days from December 8, 2019 (as initially reported by the WHO) to December 1. As a result, estimates through January 22 utilize a start date of December 8, whereas those that follow utilize a start date of December 1.

## References:

- [1] Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **2020**;DOI:[https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- [2] 疫情通报. National Health Commission of the People's Republic of China, January 2020. ([http://www.nhc.gov.cn/xcs/yqtb/list\\_gzbd.shtml](http://www.nhc.gov.cn/xcs/yqtb/list_gzbd.shtml)).
- [3] 2020 Disease Outbreak News Reports. World Health Organization, January 2020. (<https://www.who.int/csr/don/archive/year/2020/en/>).
- [4] 公示公告. Wuhan Municipal Health Commission, January 2020. (<http://wjw.wuhan.gov.cn/front/web/list2nd/no/710>).
- [5] Majumder MS, Santillana M, Mekaru SR, et al. Utilizing Nontraditional Data Sources for Near Real-Time Estimation of Transmission Dynamics During the 2015-2016 Colombian Zika Virus Disease Outbreak. *JMIR Public Health Surveill* **2016**; 2(1):e30.
- [6] Fisman DN, Hauck TS, Tuite AR, Greer AL. An IDEA for Short Term Outbreak Projection: Nearcasting Using the Basic Reproduction Number. *PLoS One* **2013**;8(12):e83622.
- [7] Majumder MS, Rivers C, Lofgren E, Fisman D. Estimation of MERS-Coronavirus Reproductive Number and Case Fatality Rate for the Spring 2014 Saudi Arabia Outbreak: Insights from Publicly Available Data. *PLoS Curr* **2014**;6:ecurrents.outbreaks.98d2f8f3382d84f390736cd5f5fe133c.
- [8] Fisman D, Khoo E, Tuite A. Early Epidemic Dynamics of the West African 2014 Ebola Outbreak: Estimates Derived with a Simple Two-Parameter Model. *PLoS Curr* **2014**;6:ecurrents.outbreaks.89c0d3783f36958d96ebbae97348d571.
- [9] Bell DM, World Health Organization Working Group on Prevention of International and Community Transmission of SARS. Public Health Interventions and SARS Spread, 2003. *Emerg Infect Dis* **2004**; 10(11):1900–1906.
- [10] Cauchemez S, Nouvellet P, Cori A, et al. Unraveling the drivers of MERS-CoV transmission. *Proc Natl Acad Sci USA* **2016**;113(32):9081–9086.



**Panel A**



**Panel B**